

Does the 2D Hubbard Model Really Show d -Wave Superconductivity?

Gang Su*

Department of Physics, Graduate School, Chinese Academy of Sciences, P.O. Box 3908, Beijing 100039, China

PACS numbers: 74.20. Mn, 74.25.Dw, 74.25.Jb

In a recent Letter [1], Maier et al concluded that they found, based on the dynamical cluster as well as non-crossing approximations on Green functions, a stable pure d -wave solution with off-diagonal long-range order (ODLRO) in the intermediate to strong coupling regime for the doped repulsive 2D Hubbard model, and given an estimation of maximum T_c being 150K for doping $x \approx 0.2$.

This is a surprising yet too strong statement. The authors in Ref. [1] might be unaware of a theorem, proved by Su and Suzuki [2] two years ago, that the 2D Hubbard model with narrow bands (including next nearest-neighbor hopping, etc.) does not exhibit d -wave pairing long-range order (LRO) at any nonzero temperature for both repulsive and attractive Coulomb interactions and for any electron fillings. This rigorous result has been generalized to cover other models (see, e.g. [3]). There are also a few rigorous results showing the absence of s -wave or other-wave pairing LROs for the 2D Hubbard model at finite temperatures [4,5]. It seems that the statement in Ref. [1] contradicts these mathematically rigorous results. On the aspect of numerical simulations, although some quantum Monte Carlo (QMC) calculations show tendencies favoring d -wave superconductivity [6], no definite sign of LRO has been detected, as pointed out by many authors (see, e.g. [7–9]). Even though the long-tailed enhancements in d -wave pairing correlations near half-filling or d -wave-like pairing fluctuations at low temperatures [6] were observed in QMC studies, none concluded from QMC results that the d -wave superconductivity with ODLRO is clearly obtained. In fact, the QMC results, owing to the well-known limitations, though still controversial, are not incompatible with the rigorous results. There indeed exist analytic and numerical works (e.g. QMC) [10–12], in completely opposite to the statement claimed in Ref. [1], showing that the 2D Hubbard model (with a next nearest-neighbor hopping integral) does not exhibit any definite sign of s -wave and d -wave superconductivity. Why are the consequences gained in Ref. [1] inconsistent with the rigorous results?

Now, let us look at what was really observed in Ref. [1]. First, the method used in Ref. [1] is only approximate. They used the coarse grained Green function to mimic the true Green function. According to the definition, as the coarse grid number N_c tends to infinity, the coarse grained Green function recovers the true Green function, whereas in the realistic presentations they use just the result of $N_c = 4$ to draw final conclusions without giving a convincing comparison for different N_c 's, apart

from just stating the systematic errors on the order of $1/N_c^3$. Second, one may notice that the statement of the appearance of a *pure* d -wave superconductivity with ODLRO comes from such an observation that the anomalous coarse grained Green function vanishes at the points $(0,0)$ and (π,π) but is finite at $(0,\pi)$ and $(\pi,0)$. Despite how reliable the approximate method is, one cannot, just from this simple observation, conclude that a *pure* d -wave superconductivity is uncovered. For example, a mixing state like $d_{x^2-y^2} + id_{xy}$ cannot be ruled out. Third, the transition observed in Ref. [1] was speculated as a possibility of a Kosterlitz-Thouless (KT) phase transition. As is well-known, the correlation function in the KT phase decays in a power law, while the existence of ODLRO implies, as discussed by Yang [13], that the correlation function becomes a nonzero constant as the spatial separation between the pairing operators tends to infinity. In their seminar article [14], Kosterlitz and Thouless also pointed clearly out that *this type of phase transition cannot occur in a superconductor*. As a result, one cannot again just from the approximate results conclude the occurrence of ODLRO in the 2D Hubbard model at low temperatures.

In short, the results presented in Ref. [1] are not sufficient to allow for drawing the conclusion like the authors did, at least in a conventional sense. As a matter of fact, it appears that the transition they observed might be a kind of topological phase transition with a power law decay (i.e. possessing a quasi-LRO), not a genuine superconducting transition because the latter possesses a true LRO. This reinterpretation makes the observation in Ref. [1] consistent with the existing rigorous results, namely, the 2D Hubbard model does not show d -wave pairing LRO at finite temperatures.

*E-mail: gsu@cc6.gsbustc.ac.cn.

- [1] Th. Maier, M. Jarrell, Th. Pruschke and J. Keller, Phys. Rev. Lett. **85**, 1524 (2000).
- [2] G. Su and M. Suzuki, Phys. Rev. B **58**, 117 (1998).
- [3] C. Noce et al, Phys. Rev. B **59** (1999), 7409.
- [4] T. Koma and H. Tasaki, Phys. Rev. Lett. **68**, 2348 (1992).
- [5] G. Su, A. Schadschneider and J. Zittartz, Phys. Lett. A **230**, 99 (1997).
- [6] D.J. Scalapino, Phys. Rep. **250**, 329 (1995); cond-mat/9606217; 9606218; 9908287.

- [7] E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).
- [8] F.F. Assaad, M. Imada and D.J. Scalapino, Phys. Rev. Lett. **77**, 4592 (1996).
- [9] K. Kuroki and H. Aoki, cond-mat/9707277.
- [10] A.F. Veilleux et al, Phys. Rev. B **52**, 16255 (1995).
- [11] T. Husslein et al, Phys. Rev. B **54**, 16179 (1996).
- [12] S. Zhang, J. Carlson and J.E. Gubernatis, Phys. Rev. Lett. **78**, 4486 (1997); Phys. Rev. B **55**, 7464 (1997).
- [13] C.N. Yang, Rev. Mod. Phys. **34**, 694 (1962). See also H.T. Nieh, G. Su and B.H. Zhao, Phys. Rev. B **51**, 3760 (1995).
- [14] J.M. Kosterlitz and D.J. Thouless, J. Phys. C **6**, 1181 (1973).